

05. Multiparticle Systems

I. 2-Particle Product Spaces

- Suppose: Particle₁ and particle₂ are represented by vector spaces V and W .
- Then: The composite 2-particle system is represented by a *product vector space* $V \otimes W$.

Let V and W be n -dim and m -dim vector spaces. The **product vector space** $V \otimes W$ is an $(n \times m)$ -dim vector space with the following property:

For any $|v\rangle \in V$, $|w\rangle \in W$, one can form a vector $|\psi\rangle \in V \otimes W$ via the "tensor product" $|v\rangle \otimes |w\rangle$, which satisfies:

- (i) $(|v_1\rangle + |v_2\rangle) \otimes |w\rangle = |v_1\rangle \otimes |w\rangle + |v_2\rangle \otimes |w\rangle$
- (ii) $|v\rangle \otimes (|w_1\rangle + |w_2\rangle) = |v\rangle \otimes |w_1\rangle + |v\rangle \otimes |w_2\rangle$
- (iii) $\alpha(|v\rangle \otimes |w\rangle) = \alpha|v\rangle \otimes |w\rangle = |v\rangle \otimes \alpha|w\rangle$, for any scalar α .

Instead of " $|v\rangle \otimes |w\rangle$ ", we can alternatively write " $|v\rangle|w\rangle$ " or " $|vw\rangle$ ".

Further characteristics of the 2-particle product space $V \otimes W$

$V \otimes W$ inherits an inner-product, bases, and operators from V and W :

1. An **inner-product** on $V \otimes W$ is defined by the following: For any $|\psi\rangle = |vw\rangle, |\phi\rangle = |tu\rangle \in V \otimes W$, with $|v\rangle, |t\rangle \in V$ and $|w\rangle, |u\rangle \in W$,
$$\langle\psi|\phi\rangle \equiv \langle v|t\rangle\langle w|u\rangle$$
2. If $\{|v_1\rangle, |v_2\rangle, \dots, |v_n\rangle\}$ and $\{|w_1\rangle, |w_2\rangle, \dots, |w_m\rangle\}$ are bases for V and W , then a **basis** for $V \otimes W$ is given by

$$\{|v_1w_1\rangle, |v_1w_2\rangle, \dots, |v_1w_m\rangle, |v_2w_1\rangle, \dots, |v_nw_m\rangle\}$$

3. Any vector $|\psi\rangle$ in $V \otimes W$ can be *expanded* in this basis:

$$|\psi\rangle = a_{11}|v_1w_1\rangle + a_{12}|v_1w_2\rangle + \dots + a_{21}|v_2w_1\rangle + \dots + a_{nm}|v_nw_m\rangle$$

4. Let A and B be operators on V and W such that

$$A|v\rangle = a|v\rangle, B|w\rangle = b|w\rangle, \quad \text{where } |v\rangle \in V, |w\rangle \in W.$$

Then there is an **operator** $A \otimes B$ on $V \otimes W$ such that

$$(A \otimes B)|vw\rangle = ab|vw\rangle$$

Extension to multiparticle (multi-partite) systems

- A product vector space \mathcal{H} may be formed from the tensor product of more than two lower-dim vector spaces.
- A product vector space \mathcal{H} may admit more than one decomposition into lower-dim vector spaces.

Ex. Let \mathcal{H} be a 16-dim vector space.

Then: One can always find 2-dim vector spaces V_1, V_2, V_3, V_4 such that

$$\mathcal{H} = V_1 \otimes V_2 \otimes V_3 \otimes V_4$$

And: One can always find 4-dim vector spaces W_1, W_2 such that

$$\mathcal{H} = W_1 \otimes W_2$$

- Note: A "factor" vector space must have $\dim > 1$.
- So: As long as the dimension n of a vector space isn't a *prime number*, it will admit at least one decomposition into the tensor product of lower-dim vector spaces.
- And: How many it will admit depends on the *prime factorization* of n .

Two-particle example


- Let: V, W be the 2-dim spin state spaces for two electrons.
 - Combined 2-particle spin space is given by 4-dim $V \otimes W$.
- Suppose:
 - $\{|hard\rangle_1, |soft\rangle_1\}$ is a basis for V .
 - $\{|hard\rangle_2, |soft\rangle_2\}$ is a basis for W .
- Then: A basis for $V \otimes W$ is given by
 - $\{|hard\rangle_1|hard\rangle_2, |hard\rangle_1|soft\rangle_2, |soft\rangle_1|hard\rangle_2, |soft\rangle_1|soft\rangle_2\}$
- And: Any 2-particle state $|A\rangle$ in $V \otimes W$ can be expanded in this basis:

$$\begin{aligned} |A\rangle = & a_{11}|hard\rangle_1|hard\rangle_2 + a_{12}|hard\rangle_1|soft\rangle_2 \\ & + a_{21}|soft\rangle_1|hard\rangle_2 + a_{22}|soft\rangle_1|soft\rangle_2 \end{aligned}$$

II. Entangled States

An **entangled state** in a product vector space \mathcal{H} with respect to a decomposition $\mathcal{H} = V_1 \otimes \cdots \otimes V_n$ is a vector $|\psi\rangle$ that *cannot* be written as a product of n terms,

$$|\psi\rangle = |v_1\rangle \otimes \cdots \otimes |v_n\rangle, \quad \text{where } |v_i\rangle \in V_i$$



Erwin Schrödinger
(1887-1961)

- What this means: An entangled n -particle state cannot be written as a product of a particle₁ state, and a particle₂ state, and a particle₃ state, *etc.*
 - *In an entangled n -particle state, the states of all n particles are "entangled with each other": they cannot be separated out.*

Two initial observations

1. Nothing about this mathematical definition tells us what the notion of "entangled with each other" means physically.
2. Entanglement is a *relative* property!
 - *A vector in \mathcal{H} can be entangled with respect to one decomposition of \mathcal{H} , but not entangled with respect to another decomposition of \mathcal{H} .*

Examples:

- Entangled: $|\Psi^+\rangle = \sqrt{1/2} \{ |hard\rangle_1 |hard\rangle_2 + |soft\rangle_1 |soft\rangle_2 \}$

- Nonentangled (Separable):

$$\begin{aligned} |A\rangle &= \sqrt{1/4} \{ |hard\rangle_1 |hard\rangle_2 + |hard\rangle_1 |soft\rangle_2 + |soft\rangle_1 |hard\rangle_2 + |soft\rangle_1 |soft\rangle_2 \} \\ &= \sqrt{1/4} \{ |hard\rangle_1 + |soft\rangle_1 \} \{ |hard\rangle_2 + |soft\rangle_2 \} \end{aligned}$$

$$|B\rangle = \sqrt{1/2} \{ |hard\rangle_1 |hard\rangle_2 + |soft\rangle_1 |hard\rangle_2 \} = \sqrt{1/2} \{ |hard\rangle_1 + |soft\rangle_1 \} |hard\rangle_2$$

$$|C\rangle = |hard\rangle_1 |hard\rangle_2$$

According to the Eigenvalue-Eigenvector Rule:

- In states $|\Psi^+\rangle$ and $|A\rangle$, both electrons have no determinate Hardness value, but the combined system *as a whole* does have a determinate value of some other property.*
- In state $|B\rangle$, electron₁ has no determinate Hardness value, but electron₂ *does* (i.e., *hard*), and the combined system as a whole has a determinate value of some other property.*
- In state $|C\rangle$, both electrons have determinate Hardness values, and the combined system as a whole has a determinate value of some other property.*

*General fact: Any vector is the eigenvector of some operator.

- So: According to the EE Rule, in any 2-particle state, either particle may or may not have well-defined properties.
- But: According to EE, *the combined 2-particle system as a whole will always have well-defined properties!*
- Why? Because, again, any vector in a vector space (including $V \otimes W$) is an eigenvector of some (Hermitian) operator on that space.
 - So there exist 2-particle operators with eigenvectors $|\Psi^+\rangle$, $|A\rangle$, $|B\rangle$ and $|C\rangle$ that represent properties of the 2-particle system *as a whole*.

2-Particle "Holistic" Properties

Suppose: $|Q\rangle = \sqrt{1/2} \{ |5\rangle_1 |7\rangle_2 + |9\rangle_1 |11\rangle_2 \}$

$|5\rangle_1$ is an eigenvector of
the position operator $X^{(1)}$

$|11\rangle_2$ is an eigenvector of
the position operator $X^{(2)}$

- If we only want to measure P1's position (and not P2's), we must use the 2-particle operator $X^{(1)} \otimes I^{(2)}$, where $I^{(2)}$ = identity operator on W .
- If we only want to measure P2's position (and not P1's), we must use the 2-particle operator $I^{(1)} \otimes X^{(2)}$, where $I^{(1)}$ = identity operator on V .
- The *difference in the positions* of P1 and P2 is a property of the 2-particle system represented by the 2-particle operator $(I^{(1)} \otimes X^{(2)}) - (X^{(1)} \otimes I^{(2)})$.

Claim: $|Q\rangle$ is an eigenstate of $(I^{(1)} \otimes X^{(2)}) - (X^{(1)} \otimes I^{(2)})$
but *not* of $X^{(1)} \otimes I^{(2)}$ or $I^{(1)} \otimes X^{(2)}$!

- So: According to the EE Rule, P1 and P2 have no definite position in the 2-particle state $|Q\rangle$, but the difference in their positions *is* a definite property of the 2-particle state as a whole!

Claim: $|Q\rangle$ is an eigenstate of $(I^{(1)} \otimes X^{(2)}) - (X^{(1)} \otimes I^{(2)})$
but *not* of $X^{(1)} \otimes I^{(2)}$ or $I^{(1)} \otimes X^{(2)}$!

Check:

(a) $|Q\rangle$ is an eigenstate of $(I^{(1)} \otimes X^{(2)}) - (X^{(1)} \otimes I^{(2)})$:

$$\begin{aligned} \{(I^{(1)} \otimes X^{(2)}) - (X^{(1)} \otimes I^{(2)})\}|Q\rangle &= (I^{(1)} \otimes X^{(2)})|Q\rangle - (X^{(1)} \otimes I^{(2)})|Q\rangle \\ &= (I^{(1)} \otimes X^{(2)})\sqrt{1/2}\{|5\rangle_1|7\rangle_2 + |9\rangle_1|11\rangle_2\} - (X^{(1)} \otimes I^{(2)})\sqrt{1/2}\{|5\rangle_1|7\rangle_2 + |9\rangle_1|11\rangle_2\} \\ &= \sqrt{1/2}\{7|5\rangle_1|7\rangle_2 + 11|9\rangle_1|11\rangle_2\} - \sqrt{1/2}\{5|5\rangle_1|7\rangle_2 + 9|9\rangle_1|11\rangle_2\} \\ &= \sqrt{1/2}\{2|5\rangle_1|7\rangle_2 + 2|9\rangle_1|11\rangle_2\} = 2|Q\rangle \end{aligned}$$

In the state represented by $|Q\rangle$, the value of the difference-in-position operator is 2; i.e., P1 and P2 differ in position by 2.

(b) $|Q\rangle$ is not an eigenstate of $X^{(1)} \otimes I^{(2)}$ or $I^{(1)} \otimes X^{(2)}$:

$$\begin{aligned} X^{(1)} \otimes I^{(2)}|Q\rangle &= (X^{(1)} \otimes I^{(2)})\sqrt{1/2}\{|5\rangle_1|7\rangle_2 + |9\rangle_1|11\rangle_2\} \\ &= \sqrt{1/2}\{5|5\rangle_1|7\rangle_2 + 9|9\rangle_1|11\rangle_2\} \\ &\neq \lambda|Q\rangle, \text{ for any value of } \lambda. \end{aligned}$$

- Similarly for $I^{(1)} \otimes X^{(2)}$.

III. Born Rule for 2-Particle States

1. Suppose a 2-particle system is in a state represented by $|k\rangle$, and suppose we measure properties of *both* P1 and P2 represented by operators $A^{(1)}$ and $B^{(2)}$.

Then: The probability that the value of $A^{(1)}$ is a_i and the value of $B^{(2)}$ is b_i is:

$$\Pr(\text{value of } A^{(1)} \text{ is } a_i \text{ and value of } B^{(2)} \text{ is } b_i \text{ in state } |k\rangle) \equiv |\langle a_i b_i | k \rangle|^2$$

where $|a_i b_i\rangle$ is an eigenvector of the 2-particle operator $A^{(1)} \otimes B^{(2)}$

2. Suppose a 2-particle system is in a state represented by $|k\rangle$, and *only* the property of P1, represented by $A^{(1)}$, is measured.

Then: The probability that the value of $A^{(1)}$ is a_i is:

$$\Pr(\text{value of } A^{(1)} \text{ is } a_i \text{ in state } |k\rangle) \equiv |\langle a_i \ell_1 | k \rangle|^2 + \cdots + |\langle a_i \ell_N | k \rangle|^2$$

where $|a_i \ell_j\rangle$, $j = 1, \dots, N$, are eigenvectors of the 2-particle operator $A^{(1)} \otimes L^{(2)}$, for any P2 property represented by $L^{(2)}$

Motivation (Law of Total Probability): The probability that the value of $A^{(1)}$ is a_i is equal to the sum of the probabilities of *all* the different ways in which the value of $A^{(1)}$ *could* be a_i .

IV. 2-Particle Projection Postulate

- Suppose: A 2-particle system is in a state represented by $|D\rangle$, and a property of P1 represented by $A^{(1)}$ is measured with the resulting value a_i .
- Then: $|D\rangle$ collapses to the state given by the following:
 - (a) Expand $|D\rangle$ in eigenvectors of the 2-particle operator $A^{(1)} \otimes L^{(2)}$, for *any* arbitrary operator $L^{(2)}$:

$$|D\rangle = d_{11}|a_1 \ell_1\rangle + \cdots + d_{1N}|a_1 \ell_N\rangle + d_{21}|a_2 \ell_1\rangle + \cdots + d_{NN}|a_N \ell_N\rangle$$

- (b) Throw out all terms other than ones with a_i . Then divide by an appropriate normalization term Λ to make sure the result is a vector with unit length:

$$|D\rangle \xrightarrow{\text{collapse}} \frac{d_{i1}|a_i \ell_1\rangle + d_{i2}|a_i \ell_2\rangle + \cdots}{\Lambda}$$

Example 1 (collapse of separable state)

- Suppose: $|D\rangle = |q_3\rangle|m_4\rangle$ is an eigenvector of $Q^{(1)} \otimes M^{(2)}$.
- Now: Suppose the property represented by $A^{(1)}$ is measured with the resulting value a_5 .
- What happens to $|D\rangle$?

- First: Expand $|D\rangle$ in the eigenvectors of $A^{(1)} \otimes L^{(2)}$, for *any* arbitrary $L^{(2)}$.
- Note: The P2 part of $|D\rangle$ already is an eigenvector of $M^{(2)}$.
- So: Use eigenvectors of $A^{(1)} \otimes M^{(2)}$ for simplicity.
$$\begin{aligned}|D\rangle &= |q_3 m_4\rangle \\ &= d_1|a_1 m_4\rangle + d_2|a_2 m_4\rangle + \cdots + d_N|a_N m_4\rangle\end{aligned}$$
- Next: Throw out all terms other than ones with a_5 , and normalize the result.
- This just leaves $d_5|a_5 m_4\rangle$.
- To normalize it, divide by its length, which is just d_5 .

• So:

$$|D\rangle \xrightarrow{\text{collapse}} |a_5\rangle|m_4\rangle$$

No change to state of P2.

Example 2 (collapse of entangled state)

- Suppose: $|D\rangle = \frac{1}{\sqrt{2}} \{|a_4 \ell_7\rangle + |a_5 \ell_{24}\rangle\}$
- Now: Suppose the property represented by $A^{(1)}$ is measured with the resulting value a_5 .
- What happens to $|D\rangle$?

Note: The P1 part of $|D\rangle$ is *already* in an eigenvector basis of $A^{(1)}$.

So: Simply throw out all terms in $|D\rangle$ that don't contain a_5 .


Result: $|D\rangle = |a_5 \ell_{24}\rangle$

• So:

$$|D\rangle \xrightarrow{\text{collapse}} |a_5\rangle |\ell_{24}\rangle$$

The state of the unmeasured P2 changes!

V. 2-Path Experiment Again.



Without Barrier:

100% of exiting electrons are white.


- At t_1 , the electron's state is: $|white\rangle|x_1, y_1\rangle = \frac{1}{\sqrt{2}} \{|hard\rangle|x_1, y_1\rangle + |soft\rangle|x_1, y_1\rangle\}$

Definite
color state

Definite
position state

- One particle with two properties in a "two-property" state.
- Represented by product vector in a product vector space.
- Just like a "2-particle" product vector for two particles, each with a single property.

V. 2-Path Experiment Again.




Without Barrier:

100% of exiting
electrons are white.

- At t_1 , the electron's state is: $|white\rangle|x_1, y_1\rangle = \frac{1}{\sqrt{2}} \{|hard\rangle|x_1, y_1\rangle + |soft\rangle|x_1, y_1\rangle\}$
 - At t_2 , the electron's state is: $\frac{1}{\sqrt{2}} \{|hard\rangle|x_2, y_2\rangle + |soft\rangle|x_3, y_1\rangle\}$
 - At t_3 , the electron's state is: $\frac{1}{\sqrt{2}} \{|hard\rangle|x_3, y_3\rangle + |soft\rangle|x_4, y_2\rangle\}$
 - At t_4 , the electron's state is: $\frac{1}{\sqrt{2}} \{|hard\rangle|x_5, y_4\rangle + |soft\rangle|x_5, y_4\rangle\} = |white\rangle|x_5, y_4\rangle$
 - $\Pr(\text{value of } C \text{ is white in state at } t_4) = |\langle white, x_5, y_4 | white, x_5, y_4 \rangle|^2 \stackrel{\checkmark}{=} 1$
- } *entangled states!*

V. 2-Path Experiment Again.



With Barrier:

50% of exiting
electrons are white,
50% are black.

- At t_4 , the electron's state is: $|k\rangle = \sqrt{1/2} \{|hard\rangle|x_5, y_4\rangle + |soft\rangle|x_3, y_1\rangle\}$
- To measure Color at t_4 , expand $|k\rangle$ in Color basis:

$$\begin{aligned}
 |k\rangle &= (\sqrt{1/2})(\sqrt{1/2})\{|black\rangle + |white\rangle\}|x_5, y_4\rangle + (\sqrt{1/2})(\sqrt{1/2})\{|black\rangle - |white\rangle\}|x_3, y_1\rangle \\
 &= 1/2|black\rangle|x_5, y_4\rangle + 1/2|black\rangle|x_3, y_1\rangle + 1/2|white\rangle|x_5, y_4\rangle - 1/2|white\rangle|x_3, y_1\rangle
 \end{aligned}$$

- $\text{Pr}(\text{value of } C \text{ is white in state } |k\rangle) = |\langle white, x_5, y_4 | k \rangle|^2 + |\langle white, x_3, y_1 | k \rangle|^2$
 $= |1/2|^2 + |1/2|^2 \stackrel{\checkmark}{=} 1/2$